

Livestock Dewormer Comparison

CATTLE

DEWORMER NAME	ACTIVE INGREDIENT	SLAUGHTER WITHDRAWAL IN DAYS	LUNGWORMS	STOMACH WORMS	INTESTINAL WORMS	BANKRUPT WORMS	NODULAR WORMS	LIVER FLUKES	TAPEWORMS	ROUNDWORMS	KIDNEY WORMS	WHIPWORMS	EYEWORMS	NASAL BOTS	LICE	MITES	HORN FLIES	GRUBS	COMMENTS	DOSAGE/BO DY WEIGHT TREATED	ADMINISTRATION *
<u>Agri-Mectin Cattle Pour-On</u>	5 mg ivermectin per ml	48	X							X					X	X		X	Do not use in female dairy cattle of breeding age.	50 ml per 1,100 lbs / 1 ltr treats 22,000 lbs.	PO
<u>Agri-Mectin Plus Clorsulon</u>	10 mg ivermectin 100 mg clorsulon per ml	49	X					X		X					X	X		X	Do not use in female dairy cattle of breeding age or in veal calves.	1 ml subcut per 110 lbs body weight.	I
<u>Bimectin Cattle Pour-On</u>	5 mg ivermectin per ml	48	X							X					X	X	X	X	Do not use in female dairy cattle of breeding age or in veal calves.	1 ml per 22 lbs / 1 ltr treats 22,000 lbs.	PO
<u>Cydectin Cattle Injectable</u>	10 mg moxidectin per ml	21	X							X					X	X		X	Do not use in female dairy cattle of breeding age, veal calves or calves less than 8 wks old.	1 ml subcut per 110 lbs / 500 ml treats 100 - 550 lb cattle.	I
<u>Cydectin Cattle Pour-On</u>	5 mg moxidectin per ml	0	X							X					X	X	X	X	No slaughter withdrawal. No milk withholding. Do not use in veal calves.	5 ml per 110 lbs / 1 ltr treats 22,000 lbs.	PO
<u>Dectomax Cattle Pour-On</u>	5 mg doramectin per ml	42	X	X	X		X								X		X		For dairy replacement heifers less than 20 mos of age and any age or class of beef cattle, including calves, pregnant	50 ml per 1,100 lbs / 1 ltr treats 22,000 lbs.	PO

<u>Safe-Guard Flaked Meal</u>	8.9 gms fenbendazole per lb	13	X	X	X	X	X													No milk withholding.	0.25 lb per 1,000 lbs for 1 day / Treats 98,000 lbs.	FM
<u>Safe-Guard Free-Choice Mineral</u>	2.24 gms fenbendazole per lb	13	X	X	X	X	X													No milk withholding.	0.1 lb per 100 lbs over 3-6 days / Treats 25,000 lbs.	FC
<u>Safe-Guard Mini Pellets</u>	8.9 gms fenbendazole per lb	13	X	X	X	X	X													No milk withholding.	0.25 lb per 1,000 lbs for 1 day / Treats 98,000 lbs.	FM
<u>Safe-Guard Paste</u>	100 mg fenbendazole per gm	8	X	X	X	X	X													No milk withholding.	290 gm - 5 gms per 220 lbs. 92 gm - 11.5 gms per 500 lbs / 290 gms treats 29 - 440 lb, 19 - 660 lb, 14 - 880 lb or 11 - 1,100 lb cattle. 92 gms treats 8 - 500 lb cattle.	O
<u>Safe-Guard Suspension</u>	100 mg fenbendazole per ml	8	X	X	X	X	X													Safe for lactating dairy cattle with no milk withholding. Do not use in veal calves.	2.3 ml per 100 lbs / 1 ltr treats 86 - 500 lb animals or 430 - 100 lb animals.	O
<u>Valbazen Suspension</u>	113.6 mg albendazole per ml	27	X	X	X			X	X											Do not use in female dairy cattle of breeding age. Do not administer to female cattle during the first 45 days of pregnancy or for 45 days after removal of bulls.	Cattle - 4 ml per 100 lbs / 1000 ml treats 50 - 500 lb cattle.	O

* Administration Types:
FC = Free Choice
FM = Feed Mix
I = Injection
O = Oral
PO = Pour-On
TD = Top Dress

GOATS

DEWORMER NAME	ACTIVE INGREDIENT	SLAUGHTER WITHDRAWAL IN DAYS	LUNGWORMS	STOMACH WORMS	INTESTINAL WORMS	BANKRUPT WORMS	NODULAR WORMS	LIVER FLUKES	TAPEWORMS	ROUNDWORMS	KIDNEY WORMS	WHIPWORMS	EYEWORMS	NASAL BOTS	LICE	MITES	HORN FLIES	GRUBS	COMMENTS	DOSAGE/BODY WEIGHT TREATED	ADMINISTRATION *
<u>Positive Pellet Goat Dewormer</u>	0.44 gms morantel tartrate per lb	30	X																Safe for all goats, including pregnant females. No milk withholding.	1 lb per 100 lbs / 6 lbs treats 600 lbs.	FM
<u>Rumatel Crumbles</u>	4.4 gms morantel tartrate per lb	30								X									No milk withholding.	0.1 lb per 100 lbs for 1 day / Treats 25,000 lbs.	FM / TD
<u>Safe-Guard Suspension</u>	100 mg fenbendazole per ml	6	X																Do not use in lactating goats.	2.9 ml per 125 lbs/ 125 ml treats 43 - 125 lb goats.	O

* Administration Types:
 FC = Free Choice
 FM = Feed Mix
 I = Injection
 O = Oral
 PO = Pour-On
 TD = Top Dress

POULTRY

DEWORMER NAME	ACTIVE INGREDIENT	SLAUGHTER WITHDRAWAL IN DAYS	ROUNDWORMS	CECAL WORMS	CAPILLARY WORMS	COMMENTS	DOSAGE	ADMINISTRATION *
<u>DuraFend</u>	2.27 gm fenbendazole per lb	0	X	X		For growing turkeys	14.5 grams per ton feed	FM
<u>Multi-Wormer Triple Action</u>	400 gm/ton hygromycin B and 4,000 gm/ton bacitracin methylene disalicylate	3	X	X	X	No withholding required for eggs	1 scoop per 1 lb feed	FM

* Administration Types:
 FC = Free Choice
 FM = Feed Mix
 I = Injection
 O = Oral
 PO = Pour-On
 TD = Top Dress

SHEEP

DEWORMER NAME	ACTIVE INGREDIENT	SLAUGHTER WITHDRAWAL IN DAYS	LUNGWORMS	INTESTINAL WORMS	LIVER FLUKES	TAPEWORMS	ROUNDWORMS	NASAL BOTS	STOMACH WORMS							COMMENTS	DOSAGE/BODY WEIGHT TREATED	ADMINISTRATION *		
									HAEMONCHUS SPP	MARSHALLAGIA SPP	OSTERTAGIA SPP	STRONGYLOIDES SPP	TELADORSAGIA SPP	TRICHOSTRONGYLUS SPP	TRICHURIS SPP					
<u>Cydectin Sheep Drench</u>	1 mg moxidectin per ml	7					X	X										Specifically formulated for use in sheep only. Do not use in female sheep providing milk for human consumption.	1 ml per 11 lbs / 1000 ml treats 100 - 110 lb sheep.	O
<u>Ivomec Drench for Sheep</u>	2400 mcg ivermectin per 3 ml	11	X				X	X	X		X	X		X	X			Formulated for use in sheep only.	3 ml per 26 lbs / 960 ml treats 83 - 100 lb sheep.	O
<u>Privermectin Drench For Sheep</u>	2400 mcg ivermectin per 3 ml	11	X				X	X	X		X	X		X	X			Formulated for use in sheep only.	3 ml per 26 lbs / 960 ml treats 83 - 100 lb sheep.	O
<u>Prohibit Soluble Drench</u>	46.8 gms levamisole HCl per pkt	3	X	X					X					X	X				Sheep - Mix 1 pkt w/1 gal water; 1 oz per 100 lbs / 1 pkt treats 128 - 100 lb sheep.	O
<u>Valbazen Suspension</u>	113.6 mg albendazole per ml	7	X	X	X	X			X	X	X			X				Do not administer to ewes during the first 30 days of pregnancy or for 30 days after removal of rams.	Sheep - 0.75 ml per 25 lbs / 1000 ml treats 64 - 50 lb sheep.	O

* Administration Types:

FC = Free Choice

FM = Feed Mix

I = Injection

O = Oral

PO = Pour-On

TD = Top Dress

SWINE

DEWORMER NAME	ACTIVE INGREDIENT	SLAUGHTER WITHDRAWAL IN DAYS	LUNGWORMS	STOMACH WORMS	INTESTINAL WORMS	BANKRUPT WORMS	NODULAR WORMS	LIVER FLUKES	TAPEWORMS	ROUNDWORMS	KIDNEY WORMS	WHIPWORMS	EYEWORMS	NASAL BOTS	LICE	MITES	HORN FLIES	GRUBS	COMMENTS	DOSAGE/BODY WEIGHT TREATED	ADMINISTRATION *
<u>Dectomax Cattle/Swine Injection</u>	10 mg doramectin per ml	24	X							X	X				X	X				Swine - 1 ml IM per 75 lbs / 500 ml treats 1000 - 38 lb pigs.	I
<u>Ivomec Cattle/Swine Injection</u>	10 mg ivermectin per ml	18	X							X					X	X		X		Swine - 1 ml subcut per 75 lbs / 500 ml treats 1000 - 38 lb pigs.	I
<u>Safe-Guard EZ-Scoop Sow Dewormer</u>	8.172 gms fenbendazole per lb	0	X	X			X			X	X	X							No slaughter withdrawal.	Mix 1 lb medicated premix for each 2,000 lbs of pork to be dewormed with a 3-12 day supply of feed / Treats 50 - 400 lb sows or 20,000 lbs pork.	F M
<u>Wazine Piperazine, 17%</u>	17 gms piperazine sulfate per 100 ml	21					X			X									Also has indications for poultry use.	1 oz per gallon drinking water / Treats 32 - 50 lb pigs.	O
<u>Wazine Piperazine, 34%</u>	34 gms piperazine sulfate per 100 ml	21					X			X									Also has indications for poultry use.	1/2 oz per gallon drinking water / Treats 512 - 50 lb pigs.	O

* Administration Types:

FC = Free Choice

FM = Feed Mix

I = Injection

O = Oral

PO = Pour-On

TD = Top Dress